

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p><b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.</b></p>					
1. REPORT DATE (DD-MM-YYYY) 02/16/2018		2. REPORT TYPE Presentation		3. DATES COVERED (From - To) 02/16/2018-02/20/2018	
4. TITLE AND SUBTITLE Secondary cutaneous amyloidosis associated with mycosis fungoides				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Cpt Kate Kimes				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg. 4430 JBSA-Lackland, TX 78236-9908 210-292-7141				8. PERFORMING ORGANIZATION REPORT NUMBER  17347	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg. 4430 JBSA-Lackland, TX 78236-9908 210-292-7141				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release. Distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <p>Secondary localized cutaneous amyloidosis is often not clinically apparent, but may be seen histologically. It is associated with several skin tumors, and has been reported with PUVA use. To date, there are three reported cases of secondary localized cutaneous amyloidosis associated with mycosis fungoides prior to any treatment. We present a case of a 39-year-old female who presented to the dermatology clinic for evaluation of facial acne. During the exam, several 5-10mm hypopigmented patches were noted on the bilateral ventral forearms. Upon questioning, the patient reported a 16 year history of these asymptomatic lesions which had been diagnosed as vitiligo; she denied prior treatments. No lymphadenopathy was noted. Under wood's lamp, the lesions did not fluoresce. The differential diagnosis included morphea, hypopigmented mycosis fungoides, post-inflammatory hypomelanosis, or idiopathic guttate hypomelanosis. Histologically, an atypical intraepidermal lymphoid infiltrate was seen with scattered foci of globular amyloid deposition in the papillary dermis and in a perivascular distribution. Immunohistochemical stains revealed predominantly CD8 positive epidermotropic lymphocytes with partial loss of CD7. T-cell gene rearrangement studies were positive for clonality.</p>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT  UU	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Clarice Longoria
					19b. TELEPHONE NUMBER (Include area code) 210-292-7141

Secondary localized cutaneous amyloidosis is often not clinically apparent, but may be seen histologically. It is associated with several skin tumors, and has been reported with PUVA use. To date, there are three reported cases of secondary localized cutaneous amyloidosis associated with mycosis fungoides prior to any treatment. We present a case of a 39-year-old female who presented to the dermatology clinic for evaluation of facial acne. During the exam, several 5-10mm hypopigmented patches were noted on the bilateral ventral forearms. Upon questioning, the patient reported a 16 year history of these asymptomatic lesions which had been diagnosed as vitiligo; she denied prior treatments. No lymphadenopathy was noted. Under wood's lamp, the lesions did not fluoresce. The differential diagnosis included morphea, hypopigmented mycosis fungoides, post-inflammatory hypomelanosis, or idiopathic guttate hypomelanosis. Histologically, an atypical intraepidermal lymphoid infiltrate was seen with scattered foci of globular amyloid deposition in the papillary dermis and in a perivenular distribution. Immunohistochemical stains revealed predominantly CD8 positive epidermotropic lymphocytes with partial loss of CD7. T-cell gene rearrangement studies were positive for clonality. The diagnosis of stage 1A mycosis fungoides with secondary cutaneous amyloidosis was made, and treatment with clobetasol was initiated. Due to the rare association of cutaneous amyloidosis with mycosis fungoides, the possibility of systemic amyloidosis was investigated with CMP, CBC, ANA, UA, UPEP, SPEP, and echocardiogram, with no significant findings. Per the patient's preference, she was also evaluated by hematology/oncology, with no changes to diagnosis or treatment. Outstanding clinical and histologic images will be presented.

"The views expressed are those of the [author(s)][presenter(s)] and do not reflect the official views or policy of the Department of Defense or it's Components"

# Secondary localized cutaneous amyloidosis associated with mycosis fungoides

Kate Kimes, D.O., CPT, USA

San Antonio Uniformed Services Education Consortium



# Disclosures

- Kate Kimes, D.O.
- Stephanie McLeish, M.D.
- Margaret Abuzeid, M.D.
- No relevant relationships with industry to disclose
- Opinions expressed are those of the authors and are not to be construed as official or as representing those of the U.S. Air Force, U.S. Army, the Department of Defense, or the United States Government

# Case presentation

- 39-year-old Fitz IV female presented to the dermatology clinic for evaluation of facial acne
- Several 5-10mm hypopigmented macules and patches on the bilateral ventral forearms



# Differential Diagnosis

- Idiopathic guttate hypomelanosis
- Post-inflammatory hypopigmentation
- Pityriasis alba
- Lichen sclerosus
- Tinea versicolor
- Hypopigmented mycosis fungoides
- Vitiligo



# Treatment

- Diagnosis: stage 1A mycosis fungoides with secondary cutaneous amyloidosis
- Labs: CMP, CBC, ANA, UA, UPEP, SPEP and echocardiogram, with no significant findings
- Treatment: topical clobetasol

# Discussion

- Associated with several skin tumors:
  - Pilomatricoma
  - Seborrheic keratosis
  - Basal cell carcinoma
  - Bowen's disease



# Discussion

- Friction/pruritus
- Apoptotic keratinocytes and cytokeratins

# References

- Bologna JL, Jorizzo JL, Schaffer JV. Dermatology 3<sup>rd</sup> edition. Philadelphia: Elsevier Saunders, 2012.
- Izumi K, Arita K, Horie K, Hoshina D, Shimizu H. Localized cutaneous amyloidosis associated with poikilodermatous mycosis fungoides. *Acta Derm Venereol* 2014; 94:225-226.
- Nam CH, Park MK, Choi MS, Hong SP, Park BC, Kim MH. Secondary cutaneous amyloidosis in a patient with mycosis fungoides. *Ann Derm* 2017; 29(1):79-82.
- Romero LS, Kantor GR, Levin MW, Vonderheid EC. Localized cutaneous amyloidosis associated with mycosis fungoides. *J. Am Acad Dermatol* 1997; 37:124-127.



Thank you!























